

Finiteness of the number of coideal subalgebras

Skryabin S.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016 American Mathematical Society. It is proved that any finite dimensional Hopf algebra which is either semisimple or cosemisimple has finitely many right coideal subalgebras. As a consequence, over an algebraically closed base field any action of a finite dimensional cosemisimple Hopf algebra on a commutative domain factors through an action of a group algebra. This extends two results of Etingof and Walton to the case where the Hopf algebra is cosemisimple, but not necessarily semisimple.

<http://dx.doi.org/10.1090/proc/13463>

References

- [1] Michel Demazure and Pierre Gabriel, Groupes algébriques. Tome I: Géométrie algébrique, généralités, groupes commutatifs (French), Masson & Cie, Éditeur, Paris; North-Holland Publishing Co., Amsterdam, 1970. Avec un appendice Corps de classes local par Michiel Hazewinkel. MR0302656
- [2] Pavel Etingof and Shlomo Gelaki, On finite-dimensional semisimple and cosemisimple Hopf algebras in positive characteristic, *Internat. Math. Res. Notices* 16 (1998), 851–864, DOI 10.1155/S107379289800052X. MR1643702
- [3] Pavel Etingof, Dmitri Nikshych, and Viktor Ostrik, On fusion categories, *Ann. of Math.* (2) 162 (2005), no. 2, 581–642, DOI 10.4007/annals.2005.162.581. MR2183279
- [4] Pavel Etingof and Chelsea Walton, Semisimple Hopf actions on commutative domains, *Adv. Math.* 251 (2014), 47–61, DOI 10.1016/j.aim.2013.10.008. MR3130334
- [5] Murray Gerstenhaber, On the deformation of rings and algebras, *Ann. of Math.* (2) 79 (1964), 59–103. MR0171807
- [6] A. Grothendieck, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV (French), *Inst. Hautes Études Sci. Publ. Math.* 32 (1967), 361 pp. MR0238860
- [7] W. J. Haboush, Reductive groups are geometrically reductive, *Ann. of Math.* (2) 102 (1975), no. 1, 67–83. MR0382294
- [8] James E. Humphreys, Linear algebraic groups, *Graduate Texts in Mathematics*, No. 21, Springer-Verlag, New York-Heidelberg, 1975. MR0396773
- [9] Richard G. Larson and David E. Radford, Finite-dimensional cosemisimple Hopf algebras in characteristic 0 are semisimple, *J. Algebra* 117 (1988), no. 2, 267–289, DOI 10.1016/0021-8693(88)90107-X. MR957441
- [10] Richard G. Larson and David E. Radford, Semisimple cosemisimple Hopf algebras, *Amer. J. Math.* 110 (1988), no. 1, 187–195, DOI 10.2307/2374545. MR926744
- [11] Akira Masuoka, Freeness of Hopf algebras over coideal subalgebras, *Comm. Algebra* 20 (1992), no. 5, 1353–1373, DOI 10.1080/00927879208824408. MR1157912
- [12] Susan Montgomery, Hopf algebras and their actions on rings, *CBMS Regional Conference Series in Mathematics*, vol. 82, published for the Conference Board of the Mathematical Sciences, Washington, DC, by the American Mathematical Society, Providence, RI, 1993. MR1243637
- [13] David Mumford and John Fogarty, Geometric invariant theory, 2nd ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]*, vol. 34, Springer-Verlag, Berlin, 1982. MR719371

- [14] Serge Skryabin, Smoothness, flatness and deformations of subalgebras, J. Pure Appl. Algebra 144 (1999), no. 1, 29–65, DOI 10.1016/S0022-4049(99)00072-9. MR1723190
- [15] Serge Skryabin, Projectivity and freeness over comodule algebras, Trans. Amer. Math. Soc. 359 (2007), no. 6, 2597–2623 (electronic), DOI 10.1090/S0002-9947-07-03979-7. MR2286047
- [16] Dragoş Ştefan, The set of types of n -dimensional semisimple and cosemisimple Hopf algebras is finite, J. Algebra 193 (1997), no. 2, 571–580, DOI 10.1006/jabr.1996.6991. MR1458803
- [17] William C. Waterhouse, Automorphism group schemes of semisimple cosemisimple Hopf algebras, Amer. J. Math. 115 (1993), no. 3, 691–698, DOI 10.2307/2375076. MR1221839